

Homework for Test #1

- 3.9 #5, 9; 11-19 odd; 45, 49
- 4.1 #5-33 odd; 55-61 odd; 67, 83
- 4.2 #7-19odd; 27-37odd; 41,43,47,53
- 4.3 #7,17,37,43,45
- 4.4 #13, 15, 23, 31

3.9 - Differentials

Recall:

For a function f that is differentiable at c , the equation of the tangent line at the point $(c, f(c))$ is given by

$$y - f(c) = f'(c)(x - c)$$

This follows from the point-slope equation $y - y_1 = m(x - x_1)$, where the slope m is the derivative $f'(x)$ evaluated at the point $(c, f(c))$.

Since c , $f(c)$, and $f'(c)$ are all constants, if we rearrange to solve for y ,

$$y = f(c) + f'(c)(x - c)$$

y is a linear function of x , called the linear approximation or tangent line approximation to the graph of $f(x)$ at $x = c$.

$$T(x) = f(c) + f'(c)(x - c)$$

For values of x close to c , values of $y = T(x)$ can be used as approximations of the values of the original function f .

Recall that the slope of the *secant line* through two points $(c, f(c))$ and $(x, f(x))$ is given by $\frac{\Delta y}{\Delta x} = \frac{f(x) - f(c)}{x - c}$, and the slope of the *tangent line* is the limit as the distance between these two points goes to zero of this expression, which we define to be the derivative.

Noting that the change in x is $\Delta x = x - c$, or $x = c + \Delta x$ and hence $f(x) = f(c + \Delta x)$, we can write this two ways:

$$f'(c) = \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} = \lim_{\Delta x \rightarrow 0} \frac{f(c + \Delta x) - f(c)}{\Delta x}$$

Actual change in y is $\Delta y = f(x) - f(c) = f(c + \Delta x) - f(c)$.

Recalling the tangent line *approximation* equation

$$T(x) = f(c) + f'(c)(x - c) = f(c) + f'(c)\Delta x$$

$$T(x) - f(c) = f'(c) \Delta x$$

We can see that change in y can be approximated by $T(x) - f(c)$, or

Approximate change in y is $\Delta y \approx f'(c)\Delta x$.

For such an approximation, Δx is denoted dx , and is called the **differential of x** . The expression $f'(x)dx$ is denoted by dy and called the **differential of y** .

$$dy = f'(x)dx$$

In many applications, the differential of y can be used as an approximation of the actual change in y , i.e. $\Delta y \approx f'(x)dx$

All of the differentiation rules can be written in **differential form**.

By definition of differentials, we have for functions (of x) u and v :

$$du = u' dx \text{ and } dv = v' dx$$

Note that rearranged, these look like $\frac{du}{dx} = u'$ and $\frac{dv}{dx} = v'$.

For example, the Product Rule becomes:

$$d[uv] = [uv]' dx = [uv' + vu'] dx = uv' dx + vu' dx = u dv + v du$$

Differential Formulas

Constant multiple: $d[cu] = c du$

Sum or difference: $d[u \pm v] = du \pm dv$

Product: $d[uv] = u dv + v du$

Quotient: $d\left[\frac{u}{v}\right] = \frac{v du - u dv}{v^2}$

3.9 #2 $f(x) = \frac{6}{x^2}$; $(2, \frac{3}{2})$ $f(2) = \frac{3}{2}$

Compare the actual function values with the tangent line approximation near 2.

$f(x) = 6x^{-2}$
 $f'(x) = -12x^{-3} = -\frac{12}{x^3}$

$f'(2) = -\frac{12}{8} = -\frac{3}{2}$

Tangent line $T(x): y = f(c) + f'(c)(x - c) = f(2) + f'(2)(x - 2)$


$T(x) = \frac{3}{2} + \frac{-3}{2}(x - 2) = \frac{3}{2} - \frac{3}{2}x + 3$

$T(x) = \frac{9}{2} - \frac{3}{2}x$

x	1.9	1.99	2	2.01	2.1
$f(x)$	1.66	1.515	1.5	1.485	1.36
$T(x)$	1.65	1.5150	1.5	1.4850	1.35

3.9 #8 $y = 1 - 2x^2 = f(x)$; $x = 0$; $\Delta x = dx = -0.1$

Compare dy and Δy for the given values of x and Δx .



$\Delta y = f(c + \Delta x) - f(c)$

$dy = f'(c)dx$

$= f(0 + (-0.1)) - f(0)$

$= f'(0) \cdot (-0.1)$

$= 1 - 2(-0.1)^2 - (1 - 2(0)^2)$


$= -4(0)(-0.1)$

$= -2(-0.1)^2$

$= 0$

$= -2(0.01)$

$= -0.02$



Find the differential dy .

$$dy = f'(x)dx$$

12. $y = 3x^{2/3}$

$$dy = 2x^{-1/3} dx$$

16. $y = \sqrt{x} + \frac{1}{\sqrt{x}} = x^{1/2} + x^{-1/2}$


$$dy = \left(\frac{1}{2}x^{-1/2} - \frac{1}{2}x^{-3/2} \right) dx$$

14. $y = \sqrt{9-x^2}$

$$= (9-x^2)^{1/2}$$

$$dy = \frac{1}{2}(9-x^2)^{-1/2} (-2x) dx$$

20. $y = \frac{\sec^2 x}{x^2+1} = (x^2+1) \cdot (\sec^2 x)' \cdot dx - (\sec^2 x) \cdot$



$$y = \frac{\sec^2 x}{x^2+1}$$

$$[(\sec x)^2]' = 2(\sec x) \cdot \sec x \tan x$$

$$dy = \frac{(x^2+1) \cdot (\sec^2 x)' dx - (\sec^2 x) (x^2+1)' dx}{(x^2+1)^2}$$

$$= \frac{(x^2+1)(2\sec^2 x \tan x dx) - (\sec^2 x)(2x dx)}{(x^2+1)^2}$$

3.9 #46

Use differentials to approximate $\sqrt[3]{26} \approx 2.96$

$$\left. \begin{array}{l} \Delta y = f(c + \Delta x) - f(c) \\ dy = f'(x)dx \\ \Delta y \approx dy \end{array} \right\} \rightarrow f(c + \Delta x) - f(c) \approx f'(x)dx$$

$$f(c + \Delta x) \approx f(c) + f'(c)dx$$

$$f(x) = \sqrt[3]{x} = x^{1/3}, \quad c = 27 \quad ; \quad \Delta x = dx = -1$$

$$f'(x) = \frac{1}{3}x^{-2/3} = \frac{1}{3\sqrt[3]{x^2}}$$

$$\sqrt[3]{26} = \sqrt[3]{27 + (-1)} \approx \sqrt[3]{27} + \frac{1}{3 \cdot (\sqrt[3]{27})^2} \cdot (-1)$$

$$= 3 - \frac{1}{27} = \frac{80}{27} \approx 2.96$$

$$\text{Recall rules of exponents: } x^{m/n} = (x^m)^{1/n} = (x^{1/n})^m \\ = \sqrt[n]{x^m} = (\sqrt[n]{x})^m$$